

Gaudino II study from the aforementioned meta-analysis.

Dusko Nežić, MD, PhD, FETCS

Aleksandar Knežević, MD, BcS

Petar Vuković, MD, BcS

“Dedinje” Cardiovascular Institute
Belgrade, Serbia

of their article was described in previously published studies.^{2,3}

Figure 1 is a copy of the illustration used in my article.

Nicholas J. Demos, MD, PA

Jersey City, New Jersey

References

1. Benedetto U, Angeloni E, Sinatra R. Reply to the editor. *J Thorac Cardiovasc Surg.* 2010;139:1669.
2. Nežić D, Knežević A, Vuković P. Does previous percutaneous coronary intervention increase the risk of graft failure in subsequent coronary surgery? *J Thorac Cardiovasc Surg.* 2010;139:1668-9.
3. Gaudino M, Cellini P, Pragliola C, Trani C, Burzotta F, Schiavoni G, et al. Arterial versus venous bypass grafts in patients with in-stent restenosis. *Circulation.* 2005;112(9 Suppl):I265-9.
4. Benedetto U, Angeloni E, Refice S, Sinatra R. Radial artery versus saphenous vein graft patency: meta-analysis of randomized controlled trials. *J Thorac Cardiovasc Surg.* 2010;139:229-31.
5. Gaudino M, Luciani N, Glieda F, Cellini C, Pragliola C, Trani C, et al. Patients with in-stent restenosis have an increased risk of mid-term venous graft failure. *Ann Thorac Surg.* 2006;82:802-5.

doi:10.1016/j.jtcvs.2010.05.045

TREATMENT OF SUBGLOTTIC STRICTURE

To the Editor:

It was a pleasure reading the article by Mercy George and colleagues in the *Journal*.¹ I would like to note that the technique for treating benign subglottic stricture on pages 414 and 415

References

1. George M, Jaquet Y, Ikonomides C, Monnier P. *J Thorac Cardiovasc Surg.* 2010;139:411-7.
2. Demos N, Timmes JJ. Management of benign tumors and obstruction of the subglottic trachea. *J Med Soc N J.* 1969;66:57-61.
3. Demos N. A method of treating benign stricture of the subglottic trachea. *Am Surg.* 1969;35:346-7.

doi:10.1016/j.jtcvs.2010.04.041

Reply to the Editor:

We thank Dr Demos for his comment on our article on the management of pediatric subglottic stenosis with glottis involvement.

The management of glotto-subglottic stenosis is challenging, and its surgical management, therefore, has evolved over the years with descriptions of the surgical technique and its modifications, along with the use of stents. The technique we have used in this large pediatric series is different from that described earlier by Demos and colleagues in 1969.^{1,2} Notably, this series is the first pediatric series of its kind. The technique combines a subglottic resection and thyro-

tracheal anastomosis with a posterior cricoid split and costal cartilage graft for glotto-subglottic stenoses. This operation was named “extended partial cricotracheal resection.” The LT-Mold prosthesis is also different in that it calibrates the supraglottic and glottic spaces in the abducted position of the vocal cords, while restoring a triangular shape to the glottis with a sharp anterior laryngeal commissure.³ All currently available stents are round or cigar shaped and do not restore a triangular glottis, especially in the case of vocal cord synechia or grade IV transglottic stenosis. The use of the LT-Mold prosthesis has significantly diminished the incidence of postoperative granulations that require repeat endoscopic/open procedures. This study was therefore a synthesis of our experience of complex pediatric glotto-subglottic stenosis that documents the evolution of the technique with an emphasis on the silicone LT-Mold.

Mercy George, MS

Philippe Monnier, MD

Department of Otorhinolaryngology

Head and Neck Surgery

University Hospital

Lausanne, Switzerland

References

1. Demos N, Timmes JJ. Management of benign tumors and obstructions of the subglottic trachea. *J Med Soc N J.* 1969;66:57-61.
2. Demos N. A method of treating benign stricture of the subglottic trachea. *Am Surg.* 1969;35:346-7.
3. Monnier P. Airway stenting with the LT Mold: experience in 30 pediatric cases. *Int J Pediatr Otorhinolaryngol.* 2007;71:1351-9.

doi:10.1016/j.jtcvs.2010.06.005




FIGURE 1. Temporary plastic obturator in upper trachea and glottis fixed with wire. Reprinted with permission.^{2,3}

THE FUTURE OF CARDIAC SURGERY

To the Editor:

What is the future of cardiac surgery? This question undoubtedly brings to mind images of fancy new technology, robots, stents, lasers, and other minimally invasive or futuristic techniques.